

**Biological Services Program
and
Division of Ecological Services**

FWS/OBS-82/10.30
SEPTEMBER 1982

REFERENCE COPY
Do Not Remove from the Library
U. S. Fish and Wildlife Service
National Wetlands Research Center
700 Cajun Dome Boulevard
Lafayette, Louisiana 70506

HABITAT SUITABILITY INDEX MODELS: BEAVER



Fish and Wildlife Service

U.S. Department of the Interior

The Biological Services Program was established within the U.S. Fish and Wildlife Service to supply scientific information and methodologies on key environmental issues that impact fish and wildlife resources and their supporting ecosystems. The mission of the program is as follows:

- To strengthen the Fish and Wildlife Service in its role as a primary source of information on national fish and wildlife resources, particularly in respect to environmental impact assessment.
- To gather, analyze, and present information that will aid decisionmakers in the identification and resolution of problems associated with major changes in land and water use.
- To provide better ecological information and evaluation for Department of the Interior development programs, such as those relating to energy development.

Information developed by the Biological Services Program is intended for use in the planning and decisionmaking process to prevent or minimize the impact of development on fish and wildlife. Research activities and technical assistance services are based on an analysis of the issues, a determination of the decisionmakers involved and their information needs, and an evaluation of the state of the art to identify information gaps and to determine priorities. This is a strategy that will ensure that the products produced and disseminated are timely and useful.

Projects have been initiated in the following areas: coal extraction and conversion; power plants; geothermal, mineral and oil shale development; water resource analysis, including stream alterations and western water allocation; coastal ecosystems and Outer Continental Shelf development; and systems inventory, including National Wetland Inventory, habitat classification and analysis, and information transfer.

The Biological Services Program consists of the Office of Biological Services in Washington, D.C., which is responsible for overall planning and management; National Teams, which provide the Program's central scientific and technical expertise and arrange for contracting biological services studies with states, universities, consulting firms, and others; Regional Staffs, who provide a link to problems at the operating level; and staffs at certain Fish and Wildlife Service research facilities, who conduct in-house research studies.

This model is designed to be used by the Division of Ecological Services in conjunction with the Habitat Evaluation Procedures.

FWS/OBS-82/10.30
September 1982

HABITAT SUITABILITY INDEX MODELS: BEAVER

by

Arthur W. Allen
Habitat Evaluation Procedures Group
Western Energy and Land Use Team
U.S. Fish and Wildlife Service
Drake Creekside Building One
2625 Redwing Road
Fort Collins, CO 80526

Western Energy and Land Use Team
Office of Biological Services
Fish and Wildlife Service
U.S. Department of the Interior
Washington, DC 20240

This report should be cited as:

Allen, A. W. 1982. Habitat suitability index models: Beaver. U.S. Dept.
Int., Fish Wildl. Serv. FWS/OBS-82/10.30. 20 pp.

PREFACE

This document is part of the Habitat Suitability Index (HSI) Model Series (FWS/OBS-82/10), which provides habitat information useful for impact assessment and habitat management. Several types of habitat information are provided. The Habitat Use Information Section is largely constrained to those data that can be used to derive quantitative relationships between key environmental variables and habitat suitability. The habitat use information provides the foundation for HSI models that follow. In addition, this same information may be useful in the development of other models more appropriate to specific assessment or evaluation needs.

The HSI Model Section documents a habitat model and information pertinent to its application. The model synthesizes the habitat use information into a framework appropriate for field application and is scaled to produce an index value between 0.0 (unsuitable habitat) and 1.0 (optimum habitat). The application information includes descriptions of the geographic ranges and seasonal application of the model, its current verification status, and a listing of model variables with recommended measurement techniques for each variable.

In essence, the model presented herein is a hypothesis of species-habitat relationships and not a statement of proven cause and effect relationships. Results of model performance tests, when available, are referenced. However, models that have demonstrated reliability in specific situations may prove unreliable in others. For this reason, feedback is encouraged from users of this model concerning improvements and other suggestions that may increase the utility and effectiveness of this habitat-based approach to fish and wildlife planning. Please send suggestions to:

Habitat Evaluation Procedures Group
Western Energy and Land Use Team
U.S. Fish and Wildlife Service
2625 Redwing Road
Ft. Collins, CO 80526

CONTENTS

| | <u>Page</u> |
|---|-------------|
| PREFACE | iii |
| ACKNOWLEDGMENTS | v |
| HABITAT USE INFORMATION | 1 |
| General | 1 |
| Food | 1 |
| Water | 3 |
| Cover | 3 |
| Reproduction | 3 |
| Interspersion | 4 |
| Special Considerations | 4 |
| HABITAT SUITABILITY INDEX (HSI) MODEL | 5 |
| Model Applicability | 5 |
| Model Description | 7 |
| Model Relationships | 10 |
| Application of the Model | 15 |
| SOURCES OF OTHER MODELS | 15 |
| REFERENCES | 18 |

ACKNOWLEDGMENTS

I gratefully acknowledge Stephen H. Jenkins and Rebecca J. Howard for their review of this habitat model. The cover of this document was illustrated by Jennifer Shoemaker. Word processing was provided by Carolyn Gulzow and Dora Ibarra.

BEAVER (Castor canadensis)

HABITAT USE INFORMATION

General

The beaver (Castor canadensis) is a large, highly specialized aquatic rodent found in the immediate vicinity of aquatic habitats (Hoffman and Pattie 1968). The species occurs in streams, ponds, and the margins of large lakes throughout North America, except for peninsula Florida, the Arctic tundra, and the southwestern deserts (Jenkins and Busher 1979). Beavers construct elaborate lodges and burrows and store food for winter use. The species is active throughout the year and is usually nocturnal in its activities. Adult beavers are nonmigratory.

Food

Beavers are generalized herbivores; however, they show strong preferences for particular plant species and size classes (Jenkins 1975; Collins 1976a; Jenkins 1979). The leaves, twigs, and bark of woody plants are eaten, as well as many species of aquatic and terrestrial herbaceous vegetation. Food preferences may vary seasonally, or from year to year, as a result of variation in the nutritional value of food sources (Jenkins 1979).

Denney (1952) summarized the food preferences of beavers throughout North America and reported that, in order of preference, beavers selected aspen (Populus tremuloides), willow (Salix spp.), cottonwood (P. balsamifera), and alder (Alnus spp.). Although several tree species have often been reported to be highly preferred foods, beavers can inhabit, and often thrive in, areas where these tree species are uncommon or absent (Jenkins 1975). Aspen and willow are considered preferred beaver foods; however, these are generally riparian tree species that may be more available for beaver foraging but are not necessarily preferred over all other deciduous tree species (Jenkins 1981). Beavers have been reported to subsist in some areas by feeding on coniferous trees, generally considered a poor quality source of food (Brenner 1962; Williams 1965). Major winter foods in North Dakota consisted principally of red osier dogwood (Cornus stolonifera), green ash (Fraxinus pennsylvanica), and willow (Hammond 1943). Rhizomes and roots of aquatic vegetation also may be an important source of winter food (Longley and Moyle 1963; Jenkins pers. comm.). The types of food species present may be less important in determining habitat quality for beavers than physiographic and hydrologic factors affecting the site (Jenkins 1981).

Tree cutting may occur during any season of the year (Jenkins 1979). However, the most intensive amount of foraging on trees or shrubs by beavers typically occurs in late fall, after green vegetation has become dessicated,

and during early spring, prior to the availability of green vegetation. Woody vegetation may be consumed immediately, although the majority of the branches and stems are hauled to a cache for storage and later use as winter food.

An adequate and accessible supply of food must be present for the establishment of a beaver colony (Slough and Sadleir 1977). The actual biomass of herbaceous vegetation will probably not limit the potential of an area to support a beaver colony (Boyce 1981). However, total biomass of winter food cache plants (woody plants) may be limiting. Low marshy areas and streams flowing in and out of lakes allow the channelization and damming of water, allowing access to, and transportation of, food materials. Steep topography prevents the establishment of a food transportation system (Williams 1965; Slough and Sadleir 1977). Trees and shrubs closest to the pond or stream periphery are generally utilized first (Brenner 1962; Rue 1964). Jenkins (1980) reported that most of the trees utilized by beaver in his Massachusetts study area were within 30 m (98.4 ft) of the water's edge. However, some foraging did extend up to 100 m (328 ft). Foraging distances of up to 200 m (656 ft) have been reported (Bradt 1938). In a California study, 90% of all cutting of woody material was within 30 m (98.4 ft) of the water's edge (Hall 1970).

Woody stems cut by beavers are usually less than 7.6 to 10.1 cm (3 to 4 inches) dbh (Bradt 1947; Hodgdon and Hunt 1953; Longley and Moyle 1963; Nixon and Ely 1969). Jenkins (1980) reported a decrease in mean stem size cut and greater selectivity for size and species with increasing distance from the water's edge. Trees of all size classes were felled close to the water's edge, while only smaller diameter trees were felled farther from the shore.

Beavers rely largely on herbaceous vegetation, or on the leaves and twigs of woody vegetation, during the summer (Bradt 1938, 1947; Brenner 1962; Longley and Moyle 1963; Brenner 1967; Aleksuk 1970; Jenkins 1981). Forbs and grasses comprised 30% of the summer diet in Wyoming (Collins 1976a). Beavers appear to prefer herbaceous vegetation over woody vegetation during all seasons of the year, if it is available (Jenkins 1981).

Aquatic vegetation, such as duck potato (Sagittaria spp.), duckweed (Lemna spp.), pondweed (Potamogeton spp.), and water weed (Elodea spp.), are preferred foods when available (Collins 1976a). Water lilies (Nymphaea spp.), with thick, fleshy rhizomes, may be used as a food source throughout the year (Jenkins 1981). If present in adequate amounts, water lily rhizomes may provide an adequate winter food source, resulting in little or no tree cutting or food caching of woody materials. Jenkins (1981) compared the rate of tree cutting by beavers adjacent to two Massachusetts ponds that contained stands of water lilies. A pond dominated by yellow water lily (N. variegatum) and white water lily (N. odorata), which have thick rhizomes, had low and constant tree cutting activity throughout the fall. Conversely, the second pond, dominated by water shield (Bassenia schreberi), which lacks thick rhizomes, had increased fall tree cutting activity by beavers. Tree cutting was particularly evident as the water shield leaves died.

Water

Beavers require a permanent supply of water and prefer a seasonably stable water level (Slough and Sadlier 1977). Beavers can usually control water depth and stability on small streams, ponds, and lakes; however, larger rivers and lakes where water depth and/or fluctuation cannot be controlled, are often partially or wholly unsuitable for the species (Murray 1961; Slough and Sadlier 1977). Beavers are absent from sizable portions of rivers in Wyoming, due to swift water and an absence of suitable dwelling sites during periods of high and low water levels (Collins 1976b).

In riverine habitats, stream gradient is the major determinant of stream morphology and the most significant factor in determining the suitability of habitat for beavers (Slough and Sadleir 1977). Retzer et al. (1956) reported that 68% of the beaver colonies recorded in Colorado were in valleys with a stream gradient of less than 6%, 28% were associated with stream gradients from 7 to 12%, and only 4% were located along streams with gradients of 13 to 14%. No beaver colonies were recorded in streams with a gradient of 15% or more. Valleys that were only as wide as the stream channel were unsuitable beaver habitat, while valleys wider than the stream channel were frequently occupied by beavers. Valley widths of 46 m (150 ft) or more were considered the most suitable. Marshes, ponds, and lakes were nearly always occupied by beavers when an adequate supply of food was available.

Cover

Lodges or burrows, or both, may be used by beavers for cover (Rue 1964). Lodges may be surrounded by water or constructed against a bank or over the entrance to a bank burrow. Water protects the lodges from predators and provides concealment for the beaver when traveling to and from food gathering areas and caches.

The lodge is the major source of escape, resting, thermal, and reproductive cover (Jenkins and Busher 1979). Mud and debarked tree stems and limbs are the major materials used in lodge construction although lesser amounts of other woody, as well as herbaceous vegetation, may be used (Rue 1964). If an unexploited food source is available, beavers will reoccupy abandoned lodges rather than build new ones (Slough and Sadleir 1977). On lakes and ponds, lodges are frequently situated in areas that provide shelter from wind, wave, and ice action. A convoluted shoreline, which prevents the buildup of large waves or provides refuge from waves, is a habitat requirement for beaver colony sites on large lakes.

Reproduction

Reproductive and cover requirements for the beaver are the same.

Interspersion

Suitable habitat for beavers must contain all of the following: (1) stable aquatic habitat providing adequate water; (2) channel gradient of less than 15%; and, (3) quality food species present in sufficient quantity (Williams 1965).

Beaver colony territories are distinct and nonoverlapping and are the fundamental unit of a beaver population (Bradt 1938). A colonized area typically contains a series of ponds of various ages, sizes, and depths (Rutherford 1964). The beavers within each colony may establish and utilize several lodges or bank burrows, or both, within their territory. During periods of low population density, the territorial boundaries of one colony may expand to include the dams and lodges of adjacent vacant colony sites (Townsend 1953). During periods of low stream flows, floodplain populations of beavers reestablished dwelling sites and territories within the main river channel in Wyoming (Collins 1976b). The average distance moved was 262 m (286 yds).

The basic composition of a beaver colony is the extended family, comprised of a monogamous pair of adults, subadults (young of the previous year), and young of the year (Svendsen 1980). Dispersal of subadults occurs during the late winter or early spring of their second year and coincides with the increased runoff from snowmelt or spring rains. Subadult beavers have been reported to disperse as far as 236 stream km (147 mi) (Hibbard 1958), although average emigration distances range from 8 to 16 stream km (5 to 10 mi) (Hodgdon and Hunt 1953; Townsend 1953; Hibbard 1958; Leege 1968).

The daily movement patterns of the beaver centers around the lodge or burrow and pond (Rutherford 1964). The density of colonies in favorable habitat ranges from 0.4 to 0.8/km² (1 to 2/mi²) (Lawrence 1954; Aleksuk 1968; Voigt et al. 1976; Bergerud and Miller 1977 cited by Jenkins and Busher 1979). The mean distance between beaver colonies in an Alaskan riverine habitat was 1.59 km (1 mi) (Boyce 1981). The closest neighbor was 0.48 km (0.3 mi) away. The size of the colony's feeding range is a function of the interaction between the availability of food and water and the colony size (Brenner 1967). The average feeding range size in Pennsylvania, excluding water, was reported to be 0.56 ha (1.4 acre). The home range of beaver in the Northwest Territory was estimated as a 0.8 km (0.5 mi) radius of the lodge (Aleksiuk 1968). The maximum foraging distance from a food cache in an Alaskan riverine habitat was approximately 800 m (874 yds) upstream, 300 m (323 yds) downstream, and 600 m (656 yds) on oxbows and sloughs (Boyce 1981).

Special Considerations

Beavers will live in close proximity to man if all habitat requirements are met (Rue 1964). However, railways, roads, and land clearing often are adjacent to waterways and may be major limiting factors affecting beaver habitat suitability (Slough and Sadleir 1977).

Transplants of beaver may be successful on strip mined land or in new impoundments where water conditions are relatively stable (Nixon and Ely 1969). Highly acidic waters, which often occur in strip mined areas, are acceptable for beaver if suitable foods are present.

HABITAT SUITABILITY INDEX (HSI) MODEL

Model Applicability

Geographic area. This HSI model was developed for application throughout the range of the beaver. However, preferred foods may vary throughout the range of the species, depending on local availability. The food component of this model assumes that woody vegetation potentially may limit the ability of an area to support beavers. Herbaceous vegetation is an important component of the summer diet of beavers and is believed to be preferred over woody vegetation during all seasons, if available. Because herbaceous vegetation is generally available throughout the year in the southern portion of the beaver's range, it may have a more important influence on the annual diet than is indicated in this model.

Season. This model has been developed to evaluate the quality of year-round habitat for the beaver.

Cover types. This model has been developed to evaluate habitat quality in the following cover types (terminology follows that of U.S. Fish and Wildlife Service 1981): Evergreen Forested Wetland (EFW); Deciduous Forested Wetland (DFW); Evergreen Scrub-Shrub Wetland (ESW); Deciduous Scrub-Shrub Wetland (DSW); Herbaceous Wetland (HW); Riverine (R); and Lacustrine (L).

Due to the foraging behavior of the beaver, the application of this model and determination of habitat units will vary by cover type. When evaluating beaver habitat in riverine, lacustrine, and wetland cover types, the model considers the area of the cover type plus a 200 m (656 ft) band of habitat on each side of the riverine channel or surrounding the water body or wetland. Figure 1 illustrates the relationship of cover types to the suggested evaluation area.

Minimum habitat area. Minimum habitat area is defined as the minimum amount of contiguous habitat that is required before an area will be occupied by a species. Information on minimum habitat area for beavers was not found in the literature. However, it is assumed that a minimum of 0.8 km (0.5 mi) of stream channel and 1.3 km² (0.5 mi²) of lake or marshland habitat must be available before these areas are suitable for colonization by beaver. If this minimum amount of habitat is not present, the HSI is assumed to be 0.0.

Cover type

Area for evaluation

Lacustrine [> 8 ha (20 acres)]

HSI determined only for area contained within 200 m (656ft) band around lake.



Lacustrine [≤ 8 ha (20 acres)]

HSI determined for area contained within 200 m band plus area of lake.



Riverine

HSI determined for area within 200 m band on both sides of river plus area of river.



Palustrine (herbaceous wetland, forested wetlands, or shrub wetlands)

HSI determined for area contained within cover type plus area within 200 m band around wetland cover type.



Figure 1. Guidelines for determining the area to be evaluated for beaver habitat suitability under various cover type conditions.

Special model considerations. Potential beaver habitat must contain a permanent source of surface water. Lakes and reservoirs that have extreme annual or seasonal fluctuations in the water level will be unsuitable habitat for beaver. Similarly, intermittent streams, or streams that have major fluctuations in discharge (e.g., high spring runoff) or a stream channel gradient of 15% or more, will have little year-round value as beaver habitat.

Assuming that there is an adequate food source available, small lakes [< 8 ha (20 acres) in surface area] are assumed to provide suitable habitat. Large lakes and reservoirs [> 8 ha (20 acres) in surface area] must have irregular shorelines (e.g., bays, coves, and inlets) in order to provide optimum habitat for the species.

Evaluation of potential beaver habitat must be centered in and around a suitable aquatic habitat. Therefore, the following factors must be taken into consideration in order to determine if this model is applicable to the habitat being evaluated:

If aquatic component of the cover
type typically has extreme changes
in water level or flow rate or
has a channel gradient exceeding
15% - - - - -

Do not continue with model;
HSI for beaver is assumed
to be 0.0.

If aquatic component of the cover
type has moderate or no fluctuation
in water level or flow rate and
channel gradient does not extend
15% - - - - -

Continue with model to
determine HSI values for
water and food.

Verification level. This model was reviewed by Stephen H. Jenkins, Ph.D., Department of Biology, University of Nevada, and Rebecca J. Howard, Research Assistant, College of Food and Natural Resources, University of Massachusetts. Improvements suggested by these reviewers were incorporated into this model.

Model Description

Overview. The HSI model for the beaver considers the quality of life requisites for the species in each cover type. Water and winter food are the only life requisites considered because the cover and reproductive needs of the species are assumed to be identical with water requirements. It also is assumed that all of the habitat requirements of the beaver can be provided within each cover type in which it occurs. Figure 2 illustrates how the HSI is related to cover types, life requisites, and specific habitat variables.

The following sections provide a written documentation of the logic and assumptions used to translate habitat information for the beaver to the variables and equations used in the HSI model. Specifically, these sections cover: (1) identification of the variables used in the model; (2) definition and justification of the suitability levels of each variable; and (3) description of the assumed relationships between variables.

Food component. Woody and herbaceous vegetation comprise the diet of the beaver. Herbaceous vegetation is a highly preferred food source throughout the year, if it is available. Woody vegetation may be consumed during any season, although its highest utilization occurs from late fall through early spring. It is assumed that woody vegetation (trees and/or shrubs) is more limiting than herbaceous vegetation in providing an adequate food source. Therefore, this model evaluates the potential of an area to provide an adequate winter food source.




Figure 2. Tree diagram illustrating the relationships of habitat variables, life requisites, and cover types to the HSI for the beaver.

Several tree and shrub species (willow, aspen, cottonwood, and alder) have often been reported to be preferred foods; however, highly preferred species may vary in different geographic regions. Although coniferous trees and shrubs may be consumed, they are a less desirable food source for beavers than are deciduous tree species. Local variations in food preference and availability should be taken into consideration when evaluating the food component of this model.

Although beavers forage at distances up to 200 m (656 ft) from water, the majority of foraging occurs within 100 m (328 ft) of the water's edge. Even though woody vegetation may be within the optimum density and size classes, it is assumed that potential food sources farther than 100 m (328 ft) from water will be of less value than woody vegetation within 100 m (328 ft). Woody vegetation in excess of 200 m (656 ft) is assumed to have no value as a potential food source.

It is assumed that a tree and/or shrub canopy closure between 40 and 60% is an indication of optimum food availability. Tree or shrub crown closures exceeding 60% are assumed to be less suitable due to the decreased accessibility of food. Extremely dense stands result in decreased mobility and the increased likelihood of cut trees hanging up in adjacent trees. To be assigned a maximum suitability value, the dbh of trees should range from 2.5 to 15.2 cm (1 to 6 inches), and shrubs should be at least 2 m (6.6 ft) tall.

The food value in a cover type is a function of the density, size class, and species composition of woody vegetation. Optimum conditions are a stand of preferred tree and/or shrub species, of medium density, less than 15.2 cm (6 inches) dbh. An adequate food source includes some trees, or shrubs, or both. The species composition of the vegetation present influences the value obtained for density and size class. Stands of highly preferred species enhance the habitat value of the site, while foods of low preference will lower the overall food value of the site. White or yellow water lilies in lacustrine cover types may be used to supplement the winter food supply. Lakes or ponds supporting these aquatic species have a higher value as winter habitat than lacustrine cover types lacking this additional food source.

Water component. Water provides cover for the feeding and reproductive activities of the beaver. A permanent and relatively stable source of water is mandatory for suitable beaver habitat.

In riverine cover types, a major change in the rate of flow or a channel gradient exceeding 15% indicate poor or unsuitable habitat. Stream channel gradients of 6% or less have optimum value as beaver habitat. Stable water levels are of optimum value as beaver habitat, while major fluctuations in the water level or flow rate decrease the value of the site. Rivers or streams that are dry during some parts of the year are assumed to be unsuitable beaver habitat.


Lacustrine habitat types less than 8 ha (20 acres) in surface area are assumed to provide suitable habitat, if an adequate food source is present. Lacustrine cover types larger than 8 ha (20 acres) in surface area must provide physical diversity (e.g., bays, coves, and inlets) in the shoreline configuration in order to provide suitable beaver habitat. It is assumed that large reservoirs or lakes that are roughly circular in shape or are comprised of extensive stretches of straight shoreline provide little shelter from wind and wave action and, therefore, have little value as beaver habitat. Variation in the water level in lacustrine cover types results in less suitable habitat quality for beavers. Lakes or ponds that are dry during portions of the year are assumed to be unsuitable beaver habitat.

All wetland cover types (e.g., herbaceous wetland and deciduous forested wetland) must have a permanent source of surface water with little or no fluctuation in order to provide suitable beaver habitat.

Model Relationships

Suitability Index (SI) graphs for habitat variables. The relationships between various conditions of habitat variables and habitat suitability for the beaver are graphically represented in this section.


| <u>Cover type</u> | <u>Variable</u> | |
|--------------------------------|-----------------|------------------------------|
| EFW,DFW, ESW,DSW, HW,R,L | V ₁ | Percent tree canopy closure. |



EFW,DFW,
ESW,DWS,
HW,R,L

V₂


Percent of trees
in 2.5 to 15.2 cm
(1 to 6 inches) dbh
size class.



EFW,DFW,
ESW,DSW,
HW,R,L

V₃


Percent shrub crown
cover.



EFW,DFW,
ESW,DSW,
HW,R,L

V₄

Average height of
shrub canopy.




EFW,DFW,
ESW,DSW,
HW,R,L

V₅

Species composition
of woody vegetation
(trees and/or shrubs)


- A) Woody vegetation dominated ($\geq 50\%$) by one or more of the following species: aspen; willow; cottonwood; or alder.
- B) Woody vegetation dominated by other deciduous species.
- C) Woody vegetation dominated by coniferous species (e.g., fir and pine).



L

V₆


Percent of lacustrine
surface dominated by
yellow and/or white
water lily.



R

V₇

Percent stream
gradient.




EFW,DFW,
ESW,DSW,
HW,R,L

V_8

Average water fluctuation on annual basis.


- A) Small fluctuations that have no effect on burrow or lodge entrances.
- B) Moderate fluctuations that affect burrow or lodge entrances.
- C) Extreme fluctuations or water absent during part of year.



L

V_9

Shoreline development factor (see variable definition in Figure 4).



Equations. In order to obtain life requisite values for the beaver, the suitability index values for appropriate variables must be combined with the use of equations. A discussion and explanation of the assumed relationships between variables was included under Model Description. The suggested equations for obtaining food and water values for the beaver are presented by cover type in Figure 3.

| <u>Life requisite</u> | <u>Cover type</u> | <u>Equation</u> |
|---------------------------|----------------------------|-------------------------|
| Winter food | DFW,EFW, DSW,ESW, HW | $\frac{a+b+c}{2.5}$ |
| Winter food | R | $\frac{b+c}{1.5}$ |
| Winter food | L | $\frac{b+c}{1.5} + V_6$ |

where: a = woody vegetation value within actual wet-land boundary. The suggested equation is:

$$[(V_1 \times V_2)^{1/2} \times V_5]^{1/2} + [(V_3 \times V_4)^{1/2} \times V_5]^{1/2}$$

b = woody vegetation value within 100 m (328 ft) from the water's edge. The suggested equation is:

$$[(V_1 \times V_2)^{1/2} \times V_5]^{1/2} + [(V_3 \times V_4)^{1/2} \times V_5]^{1/2}$$

c = woody vegetation value within 100 m (328 ft) to 200 m (656 ft) from the water's edge. The suggested equation is:

$$0.5 \left([(V_1 \times V_2)^{1/2} \times V_5]^{1/2} + [(V_3 \times V_4)^{1/2} \times V_5]^{1/2} \right)$$

| | | |
|-------|---|---------------------------------------|
| Water | R | V_7 or V_8 , whichever is lowest. |
|-------|---|---------------------------------------|

| | | |
|-------|---|--|
| Water | L | V_8 or V_9 , whichever is lowest, if lacustrine area \geq 8 ha (20 acres) in surface area. |
|-------|---|--|

V_8 , if lacustrine area is < 8 ha (20 acres) in surface area.

| | | |
|-------|------------------------|-------|
| Water | DFW,EFW, DSW,ESW,HW | V_8 |
|-------|------------------------|-------|

Figure 3. Equations for determining life requisite values by cover type for the beaver. If equation products exceed 1.0, they should be considered equal to 1.0.

HSI determination. Based on the limiting factor concept, the HSI is equal to the lowest life requisite value obtained for either food or water.

Application of the Model

Definitions of variables and suggested field measurement techniques (Hays et al. 1981) are provided in Figure 4.

SOURCES OF OTHER MODELS

Slough and Sadleir (1977) developed a land capability classification system for beaver that related habitat variables to beaver colony site density through multiple regression analysis. The model can be used for beaver population inventory because it predicts beaver colony site density.

| <u>Variable (definition)</u> | <u>Cover types</u> | <u>Suggested technique</u> |
|--|--------------------------------|---|
| V ₁ Percent tree canopy closure [the percent of the ground surface shaded by a vertical projection of the canopies of woody vegetation ≥ 5.0 m (16.5 ft) in height]. | R,L,DFW, EFW,DSW, ESW,HW | Transect, line intercept, remote sensing |
| V ₂ Percent of trees in 2.5 to 15.2 cm (1 to 6 inches) dbh size class [the percent of trees with a dbh of 2.5 to 15.2 cm (1 to 6 inches)]. | R,L,DFW, EFW,DSW, ESW,HW | Transect, quadrat, diameter tape |
| V ₃ Percent shrub crown cover [the percent of the ground surface shaded by a vertical projection of the canopies of woody vegetation < 5 m (16.5 ft) in height]. | R,L,DFW, EFW,DSW, ESW,HW | Line intercept, quadrat, remote sensing |
| V ₄ Average height of shrub canopy (the average height from the ground surface to the top of those shrubs that comprise the uppermost shrub canopy). | R,L,DFW, EFW,DSW, ESW,HW | Line intercept, quadrat, graduated rod |
| V ₅ Species composition of woody vegetation (trees and/or shrubs) (refer to model page 12). | R,L | Transect, line intercept |

Figure 4. Definitions and suggested measurement techniques of habitat variables.

| <u>Variable (definition)</u> | <u>Cover types</u> | <u>Suggested technique</u> |
|--|--------------------------------|--|
| V ₆ Percent of lacustrine surface dominated by yellow and/or white water lily [the percent of the surface dominated by yellow water lily (<i>Nymphaea variegatum</i>) and/or white water lily (<i>N. odoreta</i>)]. | L | Line intercept, remote sensing |
| V ₇ Percent stream gradient (the vertical drop in meters or feet per kilometer or mile of stream or river channel). | R | Topographic map |
| V ₈ Average water fluctuation on an annual basis (refer to model page 13). | R,L,HW, DFW,EFW, DSW,ESW | Local data |
| V ₉ Shoreline development factor (a ratio relating the relative edge of a water body to its area. To obtain a value for shoreline configuration, measure the length of the shoreline and the area of the water body. The ratio of shoreline to area is compared to that for a circle having the same area as the water body, using the following formula: | L [≥ 8 ha (20 acres)] | Remote sensing, topographic map, dot grid, map wheel |

$$DI = \frac{l}{2\sqrt{A\pi}}$$

where DI = diversity index
 l = length of shoreline
A = area of water body

A circle would have a value equal to 1.0. The greater the deviation from a circular shape, the greater the DI value will be. Values of 3 or more are assumed to be optimum for beavers).

Figure 4. (concluded).


REFERENCES

- Aleksiuk, M. 1968. Scent-mound communication, territoriality and population regulation in beaver. *J. Mammal.* 49(4):759-762.
- _____. 1970. The seasonal food regime of arctic beavers. *Ecology* 51:264-270.
- Bergerud, A. T., and D. R. Miller. 1977. Population dynamics of Newfoundland beaver. *Can. J. Zool.* 55(2):1480-1492. Cited by Jenkins and Busher 1979.
- Boyce, M. S. 1981. Habitat ecology of an unexploited population of beavers in interior Alaska. Pages 155-186 in J. A. Chapman and D. Pursley, eds. *Worldwide Furbearer Conf. Proc. Vol. I.*
- Bradt, G. W. 1938. A study of beaver colonies in Michigan. *J. Mammal.* 19:139-162.
- _____. 1947. Michigan beaver management. Mich. Dept. Conserv., Lansing. 56 pp.
- Brenner F. J. 1962. Food consumed by beavers in Crawford County, Pennsylvania. *J. Wildl. Manage.* 26(1):104-107.
- _____. 1967. Spatial and energy requirements of beaver. *Ohio J. Sci.* 67(4):242-246.
- Collins, T. C. 1976a. Population characteristics and habitat relationships of beaver in Northwest Wyoming. Ph.D. Diss., Univ. Wyoming, Laramie [Abstract only, from Diss. Abst. Int. B Sci. Eng. 37(11):5459, 1977].
- _____. 1976b. Stream flow effects on beaver populations in Grand Teton National Park. Pages 349-352 in *Proceedings of the First Conference on Scientific Research in the National Parks*, U.S. Dept. Int. Nat. Park Serv., Trans. Proc. Series 5. Vol. I.
- Denney, R. N. 1952. A summary of North American beaver management. 1946-1948. Colo. Fish Game Dept. Rep. 28, Colo. Div. Wildl. 14 pp.
- Hall, J. G. 1970. Willow and aspen in the ecology of beaver in Sagehen Creek, California. *Ecology* 41(3):484-494.
- Hammond, M. C. 1943. Beaver on the Lower Souris Refuge. *J. Wildl. Manage.* 7(3):316-321.
- Hays, R. L., C. S. Summers, and W. Seitz. 1981. Estimating wildlife habitat variables. U.S. Dept. Int., Fish and Wildl. Serv. FWS/OBS-81/47. 173 pp.
- Hibbard, E. A. 1958. Movements of beaver transplanted in North Dakota. *J. Wildl. Manage.* 22(2):209-211.

- Hodgdon, H. W., and J. H. Hunt. 1953. Beaver management in Maine. Maine Dept. Inland Fish Game, Game Div. Bull. 3. 102 pp.
- Hoffman, R. S., and D. L. Pattie. 1968. A guide to Montana mammals: identification, habitat, distribution and abundance. Univ. Montana Printing Services, Missoula. 133 pp.
- Jenkins, S. H. 1975. Food selection by beavers: a multidimensional contingency table analysis. *Oecologia* 21:157-173.
- _____. 1979. Seasonal and year-to-year differences in food selection by beavers. *Oecolog. (Berl.)* 44:112-116.
- _____. 1980. A size-distance relation in food selection by beavers. *Ecology* 61(4):740-746.
- _____. 1981. Problems, progress, and prospects in studies of food selection by beavers. Pages 559-579 in J. A. Chapman and D. Pursley, eds. *Worldwide Furbearer Conf. Proc.*, Vol. I.
- _____. Personal communication (letter dated 4 January 1982). University of Nevada, Reno, NV.
- Jenkins, S. H., and P. E. Busher. 1979. Castor canadensis. *Am. Soc. Mammal, New York. Mammalian Species* 120:1-8.
- Lawrence, W. H. 1954. Michigan beaver populations as influenced by fire and logging. Ph.D. Diss., Univ. Michigan, Ann Arbor. 219 pp. Cited by Jenkins and Busher 1979.
- Leege, T. A. 1968. Natural movements of beavers in southeastern Idaho. *J. Wildl. Manage.* 32(4):973-976.
- Longley, W. H., and J. B. Moyle. 1963. The beaver in Minnesota. *Minn. Dept. Conserv. Tech. Bull.* 6. 87 pp.
- Murray, D. F. 1961. Some factors affecting the production and harvest of beaver in the upper Tanana River Valley, Alaska. M.S. Thesis, Univ. Alaska, Anchorage. 140 pp.
- Nixon, C. M., and J. Ely. 1969. Foods eaten by a beaver colony in southeastern Ohio. *Ohio J. Sci.* 69(5):313-319.
- Retzer, J. L., H. M. Swope, J. D. Remington, and W. H. Rutherford. 1956. Suitability of physical factors for beaver management in the Rocky Mountains of Colorado. *Colo. Dept. Game, Fish and Parks, Tech. Bull.* 2:1-32.
- Rue, L. E., III. 1964. The world of the beaver. J. B. Lippincott Co., Philadelphia and New York. 155 pp.

- Rutherford, W. H. 1964. The beaver in Colorado. Colo. Dept. Game, Fish and Parks Dept., Tech. Publ. 17. 49 pp.
- Slough, B. G., and R. M. F. S. Sadleir. 1977. A land capability classification system for beaver (Castor canadensis Kuhl). Can. J. Zool. 55(8):1324-1335.
- Svendsen, G. E. 1980. Population parameters and colony composition of beaver (Castor canadensis) in southeast Ohio. Am. Midl. Nat. 104(1):47-56.
- Townsend, J. E. 1953. Beaver ecology in western Montana with special reference to movements. J. Mammal. 34(1):459-479.
- U.S. Fish and Wildlife Service. 1981. Standards for the development of habitat suitability index models. 103 ESM. U.S. Dept. Int., Fish Wildl. Serv., Div. Ecol. Serv. n.p.
- Voigt, D. R., G. B. Kolenosky, and D. H. Pimlott. 1976. Changes in summer foods of wolves in central Ontario. J. Wildl. Manage. 40(4):663-668.
- Williams, R. M. 1965. Beaver habitat and management. Idaho Wildl. Rev. 17(4):3-7.

| | | | | |
|--|--|--|----|---|
| REPORT DOCUMENTATION PAGE | | 1. REPORT NO. FWS/OBS-82/10.30 | 2. | 3. Recipient's Accession No. |
| 4. Title and Subtitle Habitat Suitability Index Models: Beaver | | | | 5. Report Date September 1982 |
| 7. Author(s) Arthur W. Allen | | | | 6. |
| 9. Performing Organization Name and Address Habitat Evaluation Procedures Group Western Energy and Land Use Team U.S. Fish and Wildlife Service Drake Creekside Building One 2625 Redwing Road Fort Collins, CO 80526 | | | | 8. Performing Organization Rept. No. |
| 12. Sponsoring Organization Name and Address Western Energy and Land Use Team Office of Biological Services Fish and Wildlife Service U.S. Department of the Interior Washington, DC 20240 | | | | 10. Project/Task/Work Unit No. |
| | | | | 11. Contract(C) or Grant(G) No. (C) (G) |
| | | | | 13. Type of Report & Period Covered |
| 15. Supplementary Notes | | | | 14. |
| 16. Abstract (Limit: 200 words) | | | | |
| <p>Habitat preferences of the beaver (<u>Castor canadensis</u>) are described in this publication, which is one of a series of Habitat Suitability Index (HSI) models. Habitat use information is presented in a synthesis of the literature on the species-habitat requirements of the beaver, followed by the development of the HSI model. The model is presented in three formats; graphic, word, and mathematical, and is designed to provide information for use in impact assessment and habitat management activities.</p> | | | | |
| 17. Document Analysis a. Descriptors | | | | |
| Rodents Animal behavior Wildlife Habit Habitability Mathematical models b. Identifiers/Open-Ended Terms <u>Castor canadensis</u> Beaver Habitat management Habitat Suitability Index (HSI) Species-habitat requirements Impact assessment c. COSATI Field/Group | | | | |
| 18. Availability Statement RELEASE UNLIMITED | | 19. Security Class (This Report) UNCLASSIFIED | | 21. No. of Pages ii - v+20pp |
| | | 20. Security Class (This Page) UNCLASSIFIED | | 22. Price |



U.S. FISH AND WILDLIFE SERVICE REGIONAL OFFICES

REGION 1

Regional Director
U.S. Fish and Wildlife Service
Lloyd Five Hundred Building, Suite 1692
500 N.E. Multnomah Street
Portland, Oregon 97232

REGION 2

Regional Director
U.S. Fish and Wildlife Service
P.O. Box 1306
Albuquerque, New Mexico 87103

REGION 3

Regional Director
U.S. Fish and Wildlife Service
Federal Building, Fort Snelling
Twin Cities, Minnesota 55111

REGION 4

Regional Director
U.S. Fish and Wildlife Service
Richard B. Russell Building
75 Spring Street, S.W.
Atlanta, Georgia 30303

REGION 5

Regional Director
U.S. Fish and Wildlife Service
One Gateway Center
Newton Corner, Massachusetts 02158

REGION 6

Regional Director
U.S. Fish and Wildlife Service
P.O. Box 25486
Denver Federal Center
Denver, Colorado 80225

REGION 7

Regional Director
U.S. Fish and Wildlife Service
1011 E. Tudor Road
Anchorage, Alaska 99503



DEPARTMENT OF THE INTERIOR U.S. FISH AND WILDLIFE SERVICE



As the Nation's principal conservation agency, the Department of the Interior has responsibility for most of our nationally owned public lands and natural resources. This includes fostering the wisest use of our land and water resources, protecting our fish and wildlife, preserving the environmental and cultural values of our national parks and historical places, and providing for the enjoyment of life through outdoor recreation. The Department assesses our energy and mineral resources and works to assure that their development is in the best interests of all our people. The Department also has a major responsibility for American Indian reservation communities and for people who live in island territories under U.S. administration.